

VERIFICATION OF TRANSLATION

Re: JAPANESE PATENT APPLICATION NO. 1999-167736

I, Hiromichi KAKEHI, of Kitahama TNK Building,  
7-1, Dosho-machi 1-chome, Chuo-ku, Osaka-shi,  
Osaka 541-0045, Japan

hereby declare that I am the translator of the  
document attached and certify that the following is  
true translation to the best of my knowledge and  
belief.

Signature of translator

Hi Kakehi

Hiromichi KAKEHI

Dated this 6th day of October, 2004

[Document Name] Specification

[Title of the Invention] Method for Determining HIV-1  
Subtype

[Claims]

5                   [Claim 1] A method for determining HIV-1  
subtypes, which comprises the steps of amplifying nucleic  
acid using as a target sequence a portion of a nucleotide  
sequence of the env gene of HIV-1 wherein at least one of  
the 5' terminal and 3' terminal nucleotide sequences is  
10 different depending on the HIV-1 subtype, and detecting  
the subtype depending on whether or not the nucleic acid  
has been amplified.

                  [Claim 2] The method according to Claim 1,  
wherein the target sequence is 100 to 2500 nucleotides  
15 long.

                  [Claim 3] The method according to Claim 1 or  
Claim 2, wherein the sequence from the 1st through 30th  
bases from the 3' terminal and/or 5' terminal of the  
target sequence is different depending on the subtype.

20                   [Claim 4] The method according to Claim 3,  
wherein the 3' terminal of the target sequence is in the  
C3 region of the env gene of HIV-1.

                  [Claim 5] The method according to Claim 4,  
wherein the 5' terminal of the target sequence is in the  
25 C2 region of the env gene of HIV-1.

[Claim 6] The method according to any of Claims 1 through 5, wherein different amplification reactions are carried out using different primer pairs, to detect different subtypes.

5           [Claim 7] The method according to Claim 6, wherein at least two different subtypes are detected by carrying out amplification reactions at least twice with different primer pairs consisting of a primer (primer 1) that includes a sequence complementary to a portion of the  
10 nucleotide sequence (nucleotide sequence 1) that differs depending on subtype in the C3 region of the env gene of HIV-1, and a primer (primer 2) that includes a sequence complementary to a portion of the nucleotide sequence (nucleotide sequence 2) of the C2 region of the env gene  
15 of HIV-1.

          [Claim 8] The method according to any of Claims 1 through 7, wherein the first amplification reaction is carried out with the first primer pair using as a target sequence a portion of a nucleotide sequence of  
20 the env gene of HIV-1, and then the second amplification reaction is carried out with the second primer pair using as a target sequence a portion of said target sequence wherein at least one of the 5' terminal and 3' terminal nucleotide sequences is different depending on the subtype,  
25 to detect the subtype depending on whether or not the

nucleic acid has been amplified by the second amplification reaction.

[Claim 9] The method according to Claim 8, wherein the second primer pair consists of a primer  
5 (primer 1) that includes a sequence complementary to a portion of the nucleotide sequence (nucleotide sequence 1) that differs depending on subtype in the C3 region of the env gene of HIV-1, and a primer (primer 2) that includes a sequence complementary to a portion of the nucleotide  
10 sequence (nucleotide sequence 2) of the C2 region of the env gene of HIV-1; and the first primer pair consists of a primer (primer 3) that includes a sequence complementary to a portion of a nucleotide sequence (nucleotide sequence 3) of the outside region of the 3' terminal of nucleotide  
15 sequence 1 of the env gene of HIV-1, and a primer (primer 4) that includes a sequence complementary to a portion of a nucleotide sequence (nucleotide sequence 4) of the outside region of the 5' terminal of nucleotide sequence 2 of the env gene of HIV-1.

20 [Claim 10] The method according to Claim 8 or Claim 9, wherein at least two subtypes are distinguished by repeating at least once, with different second primer pairs, a series of operations comprising: the first amplification reaction that is carried out with the first  
25 primer pair using as a target sequence a portion of a

nucleotide sequence of the env gene of HIV-1; the second amplification reaction that is then carried out with the second primer pair using as a target sequence a nucleotide sequence within said target sequence; to detect subtypes  
5 depending on whether or not the nucleic acid has been amplified by the second amplification reaction.

[Claim 11] The method according to Claim 10, wherein subtypes A, B, and E are distinguished by:

(a) detecting subtype A using as the first primer  
10 pair primer 12A containing nucleotide sequence GCAATAGAAAAATTCTCCTC (Sequence ID No. 5) and primer 9AE containing nucleotide sequence CACAGTACAATGCACACATG (Sequence ID No. 8), and using as the second primer pair, primer 11QA containing nucleotide sequence  
15 CTCCTGAGGGGTTAGCAAAG (Sequence ID No. 1) and primer 10 containing nucleotide sequence AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4);

(b) detecting subtype B using as the first primer pair primer 12B containing nucleotide sequence  
20 ACAGTAGAAAAATTCCCCTC (Sequence ID No. 6) and primer 9B containing nucleotide sequence CACAGTACAATGTACACATG (Sequence ID No. 9), and using as the second primer pair primer 11BB containing nucleotide sequence CTGTGCATTACAATTTCTGG (Sequence ID No. 2) and  
25 primer 10 containing nucleotide sequence

AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4); and

(c) detecting subtype E using as the first primer pair primer 12E containing nucleotide sequence GCAATAGAAAAATTCCCCTC (Sequence ID No. 7) and primer 9AE

5 containing nucleotide sequence

CACAGTACAATGCACACATG (Sequence ID No. 8), and using as the second primer pair primer 11QE containing nucleotide sequence CTCCTGAGGGTGGTTGAAAG (Sequence ID No. 3) and primer 10 containing nucleotide sequence

10 AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4).

[Claim 12] The method according to any of Claims 1 through 11, which further comprises the steps of amplifying nucleic acid using as a target sequence a portion of a nucleotide sequence of the HIV-1 genome, the  
15 nucleotide sequence being highly conserved among all subtypes, and ascertaining the presence or absence of HIV-1 depending on whether or not the nucleic acid has been amplified.

[Claim 13] A kit for determining HIV-1 subtypes,  
20 which comprises a primer pair in which a target sequence is a portion of a nucleotide sequence of the env gene of HIV-1 wherein at least one of the 5' terminal and 3' terminal nucleotides is different depending on the subtype.

25 [Detailed Description of the Invention]

[0001]

[Technical Field to Which the Invention Pertains]

The present invention relates to a method for  
determining HIV-1 subtypes, and a kit for determining HIV-  
5 1 subtypes.

[0002]

[Prior Art]

The human immunodeficiency virus (hereinafter,  
referred to as "HIV") is a virus causing acquired immune  
10 deficiency syndrome (hereinafter, referred to as "AIDS"),  
and type 1 (HIV-1) and type 2 (HIV-2) are known. Most  
cases involve HIV-1, for which various subtypes have been  
discovered.

Determining the HIV-1 subtype in infected  
15 individuals is important for assessing the reliability of  
virological test results (particularly the drug resistance  
based on genotype or the determination of plasma HIV-1 RNA  
concentration) and the route of infection. HIV-1 subtypes  
are generally determined through the sequencing of  
20 specific regions of the virus genome and phylogenetic  
analysis of the results, but these are complicated and  
expensive procedures.

[0003]

[Problem to be Solved by the Invention]

25 Thus, an object of the present invention is to

provide a simpler method for determining HIV-1 subtypes.

Another object of the present invention is to provide a kit for determining HIV-1 subtypes.

[0004]

5 [Means for Solving the Problem]

The inventors have designed various subtype-specific primers and have successfully used them to amplify nucleic acid in samples for rapid determination of HIV-1 subtypes thereby to complete the present invention.

10 Specifically, the present invention provides a method for determining HIV-1 subtypes, which comprises the steps of amplifying nucleic acid using as a target sequence a portion of a nucleotide sequence of the env gene of HIV-1 wherein at least one of the 5' terminal and  
15 3' terminal nucleotide sequences is different depending on the HIV-1 subtype, and detecting the subtype depending on whether or not the nucleic acid has been amplified. The target sequence should be 100 to 2500 base pairs in length, and preferably 150 to 500 base pairs in length. In the  
20 above method, the sequence from the 1st through 30th bases from the 3' terminal and/or 5' terminal of the target sequence should be different depending on the subtype. For example, the 3' terminal of the target sequence may be in the C3 region of the env gene of HIV-1. The 5'  
25 terminal of the target sequence may be in the C2 region of



the env gene of HIV-1. Different subtypes can be detected by different amplification reactions using different primer pairs. For example, at least two subtypes can be detected by carrying out amplification reactions at least  
5 twice with different primer pairs consisting of a primer (primer 1) that includes a sequence complementary to a portion of the nucleotide sequence (nucleotide sequence 1) that differs depending on subtype in the C3 region of the env gene of HIV-1, and a primer (primer 2) that includes a  
10 sequence complementary to a portion of the nucleotide sequence (nucleotide sequence 2) of the C2 region of the env gene of HIV-1.

[0005]

The first amplification reaction may be carried out  
15 with the first primer pair using as a target sequence a portion of a nucleotide sequence of the env gene of HIV-1, the second amplification reaction may then be carried out with the second primer pair using as a target sequence a portion of the aforementioned target sequence wherein at  
20 least one of the 5' terminal and 3' terminal nucleotide sequences is different depending on the subtype, and the subtype may be detected depending on whether or not the nucleic acid has been amplified by the second  
amplification reaction. For example, the second primer  
25 pair consists of a primer (primer 1) that includes a

sequence complementary to a portion of the nucleotide  
sequence (nucleotide sequence 1) that differs depending on  
subtype in the C3 region of the env gene of HIV-1, and a  
primer (primer 2) that includes a sequence complementary  
5 to a portion of the nucleotide sequence (nucleotide  
sequence 2) in the C2 region of the env gene of HIV-1; and  
the first primer pair may consist of a primer (primer 3)  
that includes a sequence complementary to a portion of a  
nucleotide sequence (nucleotide sequence 3) of the outside  
10 region of the 3' terminal of nucleotide sequence 1 of the  
env gene of HIV-1, and a primer (primer 4) that includes a  
sequence complementary to a portion of the nucleotide  
sequence (nucleotide sequence 4) of the outside region of  
the 5' terminal of nucleotide sequence 2 of the env gene  
15 of HIV-1. At least two subtypes can be distinguished by  
repeating at least once the following series of operations  
with different second primer pairs, the operations  
comprising the first amplification reaction that is  
carried out with the first primer pair using as a target  
20 sequence a portion of a nucleotide sequence of the env  
gene of HIV-1, the second amplification reaction that is  
then carried out with the second primer pair using as a  
target sequence a nucleotide sequence within the above  
target sequence, and the detection of subtypes depending  
25 on whether or not the nucleic acid has been amplified by

the second amplification reaction. For example, subtypes A, B, and E can be distinguished by: (a) detecting subtype A using as the first primer pair primer 12A containing nucleotide sequence GCAATAGAAAAATTCTCCTC (Sequence ID No. 5) and primer 9AE containing nucleotide sequence CACAGTACAATGCACACATG (Sequence ID No. 8), and using as the second primer pair primer 11QA containing nucleotide sequence CTCCTGAGGGGTTAGCAAAG (Sequence ID No. 1) and primer 10 containing nucleotide sequence AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4);

(b) detecting subtype B using as the first primer pair primer 12B containing nucleotide sequence ACAGTAGAAAAATTCCCCTC (Sequence ID No. 6) and primer 9B containing nucleotide sequence CACAGTACAATGTACACATG (Sequence ID No. 9), and using as the second primer pair primer 11BB containing nucleotide sequence CTGTGCATTACAATTTCTGG (Sequence ID No. 2) and primer 10 containing nucleotide sequence AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4); and

(c) detecting subtype E using as the first primer pair primer 12E containing nucleotide sequence GCAATAGAAAAATTCCCCTC (Sequence ID No. 7) and primer 9AE containing nucleotide sequence CACAGTACAATGCACACATG (Sequence ID No. 8), and using as the second primer pair primer 11QE containing nucleotide sequence

CTCCTGAGGGTGGTTGAAAG (Sequence ID No. 3) and primer 10  
containing nucleotide sequence AAATGGCAGTCTAGCAGAAG  
(Sequence ID No. 4).

[0006]

5           The method of the present invention may further  
comprise the steps of amplifying nucleic acid using as a  
target sequence a portion of a nucleotide sequence of the  
HIV-1 genome, the nucleotide sequence being highly  
conserved among all subtypes and ascertaining the presence  
10 or absence of HIV-1 depending on whether or not the  
nucleic acid has been amplified.

Another object of the present invention is to  
provide a kit for determining HIV-1 subtypes, comprising a  
primer pair in which a target sequence is a portion of a  
15 nucleotide sequence of the env gene of HIV-1 wherein at  
least one of the 5' terminal and 3' terminal nucleotide  
sequences is different depending on the subtype.

[0007]

[Mode for Carrying out the Invention]

20           Hereinafter, embodiments of the present invention  
are illustrated below.

A sample of blood, lymph, spinal fluid, semen, lymph  
node, or the like is taken from individuals suspected of  
HIV-1 infection, infected individuals and patients  
25 confirmed with HIV-1 infection, patients being treated for

HIV-1, and the like. DNA is extracted using a QIAamp  
Blood Kit (QIAGEN), either directly or after monocytes  
have been isolated from the sample by Ficoll-Paque density  
gradient centrifugation (Pharmacia). Alternatively, RNA  
5 is extracted using a QIAamp Viral RNA Kit (QIAGEN) from  
plasma. The DNA or RNA concentration is then determined  
based on the absorption at the wavelength of 260 nm.

[0008]

The nucleic acid is then treated in PCR, and  
10 preferably nested PCR.

The use of nested PCR is described below. Nested  
PCR involves designing the second primer pair inside a  
target sequence amplified with another primer pair (first  
primer pair), carrying out the first PCR step, and then  
15 diluting the reaction product as a new template for the  
second PCR step, and then carrying out the second PCR step.  
Undesirable sequences are sometimes amplified in addition  
to the target sequence in the first PCR step. However,  
there is very little possibility that undesirable  
20 fragments amplified during the first PCR step have a  
sequence with which the primers of the second primer pair  
will anneal. The second PCR step is thus carried out for  
selectively amplifying the target sequence.

[0009]

25 The initial PCR step (first PCR) is first carried

out using different primer pairs specific to each subtype  
to be distinguished (such as subtype A, subtype B, and  
subtype E). Alternatively, universal primer pairs  
allowing any type of subtypes to be amplified can be used  
5 instead of subtype-specific primer pairs.

An example of a subtype-specific primer pair is a  
primer pair consisting of a primer (primer 4') which  
includes a sequence complementary to a portion of a  
nucleotide sequence in the C2 region of the env gene of  
10 HIV-1 and a primer (primer 3') which includes a sequence  
complementary to a portion of the nucleotide sequence of  
the C3 region of the env gene for HIV-1 that differs  
depending on subtype (that is, subtype-specific nucleotide  
sequence). Since the C2 region of the env gene of HIV-1  
15 has a nucleotide sequence that differs depending on the  
subtype, as shown in Figure 1, the nucleotide sequence may  
be selected from this region to design primer 4'. Because  
the C3 region of the env gene of HIV-1 varies depending on  
the subtype, as shown in Figure 2, the nucleotide sequence  
20 may be selected from this region to design primer 3'. The  
primer should generally be 18 to 30 base pairs, and  
preferably 20 to 25 base pairs in length. Specifically,  
the following primers can be used.

[0010]

25 A subtype A-specific primer pair for the first PCR

and their nucleotide sequences

9AE/12A

primer 9AE: CACAGTACAATGCACACATG (Sequence ID No. 8)

primer 12A: GCAATAGAAAAATTCTCCTC (Sequence ID No. 5)

5 [0011]

A subtype B-specific primer pair for the first PCR  
and their nucleotide sequences

9B/12B

primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

10 primer 12B: ACAGTAGAAAAATTCCCCTC (Sequence ID No. 6)

[0012]

A subtype E-specific primer pair for the first PCR  
and their nucleotide sequences

9AE/12E

15 primer 9AE: CACAGTACAATGCACACATG (Sequence ID No. 8)

primer 12E: GCAATAGAAAAATTCCCCTC (Sequence ID No. 7)

[0013]

A subtype C-specific primer pair for the first PCR  
and their nucleotide sequences

20 9B/12A

primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

primer 12A: GCAATAGAAAAATTCTCCTC (Sequence ID No. 5)

[0014]

A subtype D-specific primer pair for the first PCR  
25 and their nucleotide sequences

9B/12B

primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

primer 12B: ACAGTAGAAAAATTCCCCTC (Sequence ID No. 6)

[0015]

5        A subtype F-specific primer pair for the first PCR  
and their nucleotide sequences

9B/12A

primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

primer 12A: GCAATAGAAAAATTCTCCTC (Sequence ID No. 5)

10       [0016]

A subtype G-specific primer pair for the first PCR  
and their nucleotide sequences

9B/12A

primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

15       primer 12A: GCAATAGAAAAATTCTCCTC (Sequence ID No. 5)

[0017]

A subtype H-specific primer pair for the first PCR  
and their nucleotide sequences

9B/12A

20       primer 9B: CACAGTACAATGTACACATG (Sequence ID No. 9)

primer 12A: GCAATAGAAAAATTCTCCTC (Sequence ID No. 5)

[0018]

Any subtype may be amplified by using a mixture of  
the above-mentioned subtype-specific primers instead of a  
25       single use of the above subtype-specific primer pair.



1/1000 to 1/5 (for example, 1/50) of the PCR products is used to carry out the next PCR (second PCR) with another primer pair specific to each subtype. The primer pair for the second PCR is designed within the target sequence amplified during the first PCR. For example, at least one of the primers forming the subtype-specific primer pair for the second PCR can be a primer (primer 1) containing a sequence complementary to a portion of the subtype-specific nucleotide sequence of the C2 region of the env gene for HIV-1. Since the nucleotide sequence of the C2 region of the env gene for HIV-1 differs depending on subtype, as shown in Figure 1, a nucleotide sequence from this region can be selected to design the primer. Figure 2 gives the nucleotide sequence of the 3' adjacent region (C3 region) of the V3 region of the env gene for various subtypes of HIV-1. Since the nucleotide sequence varies depending on the subtype, a suitable sequence can be selected to design a primer. To design a subtype-specific primer, phylogenetic analysis is employed to select nucleotide sequences of a given subtype that are as genetically remote as possible from the corresponding nucleotide sequences of other subtypes. An example can include a primer (primer 2) containing a sequence complementary to a portion of the nucleotide sequence of the C3 region of the

env gene for HIV-1. Specifically, the primer pair containing the following nucleotide sequences can be used.

[0019]

A subtype A-specific primer pair for the second PCR  
5 and their nucleotide sequences

10/11QA

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

primer 11QA: CTCCTGAGGGGTTAGCAAAG (Sequence ID No. 1)

[0020]

10 A subtype B-specific primer pair for the second PCR  
and their nucleotide sequences

10/11BB

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

primer 11BB: CTGTGCATTACAATTTCTGG (Sequence ID No. 2)

15 [0021]

A subtype E-specific primer pair for the second PCR  
and their nucleotide sequences

10/11QE

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

20 primer 11QE: CTCCTGAGGGTGGTTGAAAG (Sequence ID No. 3)

[0022]

A subtype C-specific primer pair for the second PCR

10C/11RC

primer 10C: AAATGGTAGCCTAGCAGAAG (Sequence ID No. 10)

25 primer 11RC: CTCCTGAGGATGGTGCAAATTT (Sequence ID No. 13)

[0023]

A subtype D-specific primer pair for the second PCR  
10/11RD

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

5 primer 11RD: CTCCTGAGGATGGTTTAAAAAT (Sequence ID No. 14)

[0024]

A subtype F-specific primer pair for the second PCR  
10C/11RF

primer 10C: AAATGGTAGCCTAGCAGAAG (Sequence ID No. 10)

10 primer 11RF: CTCCTGAGGATGAGTTAAATTT (Sequence ID No. 15)

[0025]

A subtype G-specific primer pair for the second PCR  
10G/11SG

primer 10G: GAATGGCAGTTTAGCAGAAG (Sequence ID No. 11)

15 primer 11SG: TCCTGCAGATGAGTTAAAGG (Sequence ID No. 16)

[0026]

A subtype H-specific primer pair for the second PCR  
10H/11SH

primer 10H: GTCAAATGGCAGTTTAGCAG (Sequence ID No. 12)

20 primer 11SH: TCCTGAGGATGGTTTAAAGG (Sequence ID No. 17)

[0027]

A subtype universal primer pair for the second PCR  
10/11LB

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

25 primer 11LB: AATTTCTGGGTCCCCTCCTG (Sequence ID No. 18)

[0028]

A subtype universal primer pair for the second PCR  
10/11LE

primer 10: AAATGGCAGTCTAGCAGAAG (Sequence ID No. 4)

5 primer 11LE: AATTTCTAGATCTCCTCCTG (Sequence ID No. 19)

[0029]

The PCR procedures and reaction conditions may be in  
accordance with those in Bruisten S. et al., *AIDS Res Hum*  
*Retroviruses* 1993, 9:259-265, but the hot start method is  
10 preferred. In hot start PCR, the PCR reaction solution is  
kept on a hot plate for start up at an elevated  
temperature (usually 90°C or higher).

However, it sometimes happens that no subtype is  
detected in an attempt to determine the HIV-1 subtype in  
15 such a method. Possible causes may be that the HIV-1 DNA  
concentration is below the detection threshold, or the  
presence of numerous variants at the primer binding site.

[0030]

To deal with the former possibility, the above  
20 method can be implemented after extracting RNA from plasma  
and converting the RNA into cDNA using reverse  
transcriptase, since the concentration of HIV-1 is  
generally higher in plasma than in cells.

To deal with the latter possibility, the  
25 determination of the subtype by the method of the present

invention is held off, another genetic region of HIV-1 is amplified by PCR to determine the nucleotide sequence, and the subtype is determined by a conventional method (Note: HIV-1 infection is generally diagnosed by detecting  
5 antibodies. This invention is not a method for diagnosing HIV-1 infection.).

The PCR reaction products are separated by agarose gel electrophoresis and detected by ethidium bromide staining. Although distinct bands can be observed with  
10 the use of primers consistent with the subtype of the HIV-1 in sample DNA, the bands are indistinct or not observed at all when the primers are not consistent with the subtype of the HIV-1 in sample DNA. The HIV-1 subtype is determined in this way.

15 [0031]

The present invention also encompasses a kit for determining HIV-1 subtypes, comprising a primer pair in which a target sequence is a portion of a nucleotide sequence of the env gene of HIV-1 wherein at least one of  
20 the 5' terminal and 3' terminal nucleotide sequences is different depending on the subtype. Examples of primer pairs include primer pairs (inner primers) for the second PCR such as those above, and combinations of primer pairs for the first PCR (outer primers) and primer pairs for the  
25 second PCR. The kit of the present invention may also

include dNTP mixtures, reaction buffers, DNA polymerase.  
To minimize the effects caused by inconsistencies between  
the primer and analyte HIV-1 DNA base pairs, the magnesium  
ion concentration in the reaction buffer should be  
5 increased from the usual concentration of 1.5 mM to 4 mM.

The components constituting the diagnostic kit may  
be packaged individually, assembled, or bundled in  
containers such as vials and tubes, and further the  
containers may be in supporting means divided for housing  
10 such components.

[0032]

[Examples]

The present invention is illustrated in detail with  
the following examples, but the scope of the present  
15 invention is not limited to these examples.

[Example 1]

Subjects and Method

1) Subtype-specific specimens to study method for  
determining subtype

20 Specimens were prepared by extracting DNA from the  
blood of 3 HIV-infected subjects determined to be subtype  
A, 8 HIV-infected subjects determined to be subtype B and  
3 HIV-infected subjects determined to be subtype E,  
through env gene sequencing and phylogenetic analysis.

25 [0033]

2) Subjects for determining subtype

The HIV-1 subtype was determined using 8 HIV-infected subjects who either visited or were hospitalized in hospitals in Tokyo.

5 3) Preparation of DNA from blood of HIV patients

10 ml-peripheral blood was collected from the above HIV-infected patients. Sodium citrate was used as an anticoagulant. Monocytes were separated from the peripheral blood by Ficoll-Paque (Pharmacia) density  
10 gradient centrifugation, and DNA was then prepared using a QIAamp Blood Kit (QIAGEN). The DNA was dissolved in pure water or buffer containing 1 mM EDTA, and was stored at -20°C until immediately before use. 0.5 µg DNA was used in PCR.

15 [0034]

4) Detection of subtypes A, B, and E by PCR

Figure 3 gives the nucleotide sequences of the primers used in PCR.

For subtype A-specific detection, nested PCR was  
20 carried out using 9AE and 12A as the primers for the first PCR, and 10 and 11QA as the primers for the second PCR. For subtype B-specific detection, nested PCR was carried out using 9B and 12B as the primers for the first PCR, and 10 and 11BB as the primers for the second PCR. For  
25 subtype E-specific detection, nested PCR was carried out

using 9AE and 12E as the primers for the first PCR, and 10 and 11QE as the primers for the second PCR (Figure 3).

[0035]

PCR was carried out for 30 cycles, wherein one cycle  
5 consisted of 15 seconds at 94°C, 30 seconds at 56°C, and 1  
minute at 72°C, with 100 µL reaction solution (10 mM Tris-  
HCl pH 8.3, 50 mM KCl, 4 mM MgCl<sub>2</sub>, 0.2 mM dNTP, 1.0 µM  
primer, 2.5 units Taq polymerase) using 0.5 µg sample DNA  
prepared from HIV-infected subjects. Using 2 µL reaction  
10 solution obtained from the first PCR, the second PCR was  
carried out for 25 cycles under the same conditions with  
the exception of using 30 seconds at 60°C instead of 30  
seconds at 56°C.

PCR products (subtypes A and E: 317 bp; subtype B:  
15 342 bp) were detected by isolation through electrophoresis  
with 2% agarose gel, and by subsequent ethidium bromide  
staining.

[0036]

#### Results

20 1) Study of subtype determination by PCR of subtype-  
specific specimens

The following results were obtained for specimens  
whose subtypes had been determined by virus genome  
sequencing. Only subtype A specimens were positive, while  
25 subtype B and E specimens were all negative, in PCR using



primers for the detection of subtype A. Only subtype B specimens were positive, while subtype A and E specimens were all negative, in PCR using primers for the detection of subtype B. Only subtype E specimens were positive, while subtype A and B specimens were all negative, in PCR using primers for the detection of subtype E (Figure 4).

[0037]

2) Determination of subtype of HIV patients by PCR

Table 1 gives the results obtained in the determination of HIV-1 subtypes in 8 HIV-infected patients who either visited or were hospitalized in hospitals in Tokyo.

[0038]

Table 1 The results obtained in the determination of unknown subtypes in 8 specimens

| Case | primer pair for<br>subtype A | primer pair for<br>subtype B | primer pair for<br>subtype E |
|------|------------------------------|------------------------------|------------------------------|
| P18  | —                            | +                            | —                            |
| P19  | —                            | +                            | —                            |
| P20  | —                            | +                            | —                            |
| P21  | —                            | —                            | +                            |
| P22  | —                            | ?                            | —                            |
| P23  | —                            | —                            | +                            |
| P24  | —                            | +                            | —                            |
| P25  | —                            | +                            | —                            |

In the table above, + denotes detection of HIV-1 specific DNA bands, - denotes non-detection thereof. The symbol "?" for Case 22 denotes detection of shorter bands than expectation.

[0039]

Based on these results, Cases P18, P19, P20, P24, and P25 were diagnosed as being infected with subtype B, and Cases P21 and P23 were diagnosed as being infected  
5 with subtype E. Although a DNA band was detected only with the use of a primer pair for subtype B in Case P22, it was shorter than expected, so determination was postponed. To verify that the above results were correct, the amplified DNA was sequenced and phylogenetically  
10 analyzed, showing that the results of the phylogenetic analysis were consistent with those in Table 1. Case P22, which was postponed, turned out to be subtype B.

[0040]

The results in 1) and 2) demonstrate that this  
15 method was able to correctly diagnose the subtype in 21 out of 22 cases. The determination was postponed in the remaining one case. That is, the present method has been shown to be a simple and reliable method for determining subtypes.

20 The method of the present invention allows the determination of HIV-1 subtype at a cost of about ¥2,000 per specimen. The time needed to determine the subtype in treating all 8 specimens at once was 2 hours for the isolation of DNA, 6 hours for PCR, and about 1 hour for  
25 electrophoresis.

[0041]

[Effect of the Invention]

The present invention provides a simple method for  
determining HIV-1 subtypes. The invention also provides  
5 an effective means for determining HIV-1 subtypes.

[0042]

[Sequence Listing]

<110> KEIO UNIVERSITY

10 <120> A Method For HIV-1 Subtyping

<130> 17 sequences

<140>

15 <141>

<160> 19

<170> PatentIn Ver.2.0

20

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

25

$\langle 220 \rangle$ 

<223> Description of Artificial Sequence:synthetic DNA

 $\langle 400 \rangle$  1

5    ctcttgaggg gtttagcaaag

20

 $\langle 210 \rangle \quad 2$ 

$\langle 211 \rangle$  20

10 <212> DNA

### <213> Artificial Sequence

 $\langle 220 \rangle$ 

<223> Description of Artificial Sequence:synthetic DNA

15

 $\langle 400 \rangle_2$ 

ctgtgcatta caatttctgg 20

 $\langle 210 \rangle \quad 3$ 

20 <211> 20

<212> DNA

### <213> Artificial Sequence

 $\langle 220 \rangle$ 

25 <223> Description of Artificial Sequence:synthetic DNA

<400> 3

ctcctgaggg tggttgaaag

20

5 <210> 4

<211> DNA

<212> DNA

<213> Artificial Sequence

10 <220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 4

aaatggcagt ctagcagaag

20

15

<210> 5

<211> 20

<212> DNA

<213> Artificial Sequence

20

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 5

25 gcaatagaaa aattctcctc

20

<210> 6

<211> 20

<212> DNA

5 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

10 <400> 6

acagtagaaa aattcccctc 20

<210> 7

<211> 20

15 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:synthetic DNA

20

<400> 7

gcaatagaaa aattcccctc 20

<210> 8

25 <211> 20

<212> DNA

<213> Artificial Sequence

<220>

5 <223> Description of Artificial Sequence:synthetic DNA

<400> 8

cacagtacaa tgcacacatg 20

10 <210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 9

cacagtacaa tgtacacatg 20

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

25

<220>

<220>

<223> Description of Artificial Sequence:synthetic DNA

5

<400> 10

aaatggtagc ctagcagaag 20

<210> 11

10 <211> 20

<212> DNA

<213> Artificial Sequence

<220>

15

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 11

20 gaatggcagt ttagcagaag 20

<210> 12

<211> 20

<212> DNA

25 <213> Artificial Sequence



<220>

<220>

5 <223> Description of Artificial Sequence:synthetic DNA

<400> 12

gtcaaattggc agtttagcag

20

10 <210> 13

<211> 22

<212> DNA

<213> Artificial Sequence

15 <220>

<220>

<223> Description of Artificial Sequence:synthetic DNA

20 <400> 13

ctcctgagga tgggtgcaaattt

22

<210> 14

<211> 22

25 <212> DNA

<213> Artificial Sequence

<220>

5 <220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 14

ctcctgagga tggtttaaaa at

22

10

<210> 15

<211> 22

<212> DNA

<213> Artificial Sequence

15

<220>

<220>

<223> Description of Artificial Sequence:synthetic DNA

20

<400> 15

ctcctgagga tgagttaa at tt

22

<210> 16

25 <211> 20

<212> DNA

<213> Artificial Sequence

<220>

5

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 16

10 tcctgcagat gagttaaagg

20

<210> 17

<211> 20

<212> DNA

15 <213> Artificial Sequence

<220>

<220>

20 <223> Description of Artificial Sequence:synthetic DNA

<400> 17

tcctgaggat ggtttaaagg

20

25 <210> 18

<211> 20

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 18

aatttctggg tcccctcctg 20

10

<210> 19

<211> 20

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence:synthetic DNA

<400> 19

20 aatttctaga tctcctcctg 20

[0043]

[Free Text of Sequence Listing]

Sequence ID No. 1 gives the nucleotide sequence for  
primer 11QA.

25 Sequence ID No. 2 gives the nucleotide sequence for

primer 11BB.

Sequence ID No. 3 gives the nucleotide sequence for  
primer 11QE.

Sequence ID No. 4 gives the nucleotide sequence for  
5 primer 10.

Sequence ID No. 5 gives the nucleotide sequence for  
primer 12A.

[0044]

Sequence ID No. 6 gives the nucleotide sequence for  
10 primer 12B.

Sequence ID No. 7 gives the nucleotide sequence for  
primer 12E.

Sequence ID No. 8 gives the nucleotide sequence for  
primer 9AE.

Sequence ID No. 9 gives the nucleotide sequence for  
15 primer 9B.

Sequence ID No. 10 gives the nucleotide sequence for  
primer 10C.

[0045]

Sequence ID No. 11 gives the nucleotide sequence for  
20 primer 10G.

Sequence ID No. 12 gives the nucleotide sequence for  
primer 10H.

Sequence ID No. 13 gives the nucleotide sequence for  
25 primer 11RC.

Sequence ID No. 14 gives the nucleotide sequence for primer 11RD.

Sequence ID No. 15 gives the nucleotide sequence for primer 11RF.

5 [0046]

Sequence ID No. 16 gives the nucleotide sequence for primer 11SG.

Sequence ID No. 17 gives the nucleotide sequence for primer 11SH.

10 Sequence ID No. 18 gives the nucleotide sequence for primer 11LB.

Sequence ID No. 19 gives the nucleotide sequence for primer 11LE.

[Brief Description of the Drawing]

15 [Figure 1]

Figure 1 illustrates nucleotide sequences of the 5' adjacent region (C2 region) of the V3 region for the env gene in various subtypes of HIV-1. Capital letters indicate that the nucleotides are entirely the same within a given subtype, and lower case letters indicate the presence of nucleotide variants that are different within a given subtype. A question mark "?" indicates that a consensus nucleotide was not determined because of too many variants. A dash "-" indicates a nucleotide identical to that in subtype A. A period "." indicates

20

25

the absence of a nucleotide in the corresponding site.

[Figure 2]

Figure 2 illustrates nucleotide sequences of the 3' adjacent region (C3 region) of the V3 region for the env gene in various subtypes of HIV-1. Capital letters indicate that the nucleotides are entirely the same within a given subtype, and lower case letters indicate the presence of nucleotide variants that are different within a given subtype. A question mark "?" indicates that a consensus nucleotide was not determined because of too many variants. A dash "-" indicates a nucleotide identical to that in subtype A. A period "." indicates the absence of a nucleotide in the corresponding site.

[Figure 3]

Figure 3 illustrates the design of primers used in nested PCR for determining HIV-1 subtypes.

[Figure 4]

Figure 4 gives the results obtained when subtypes were detected by nested PCR for specimens in which the subtypes had been determined by sequencing of the virus genome.

[Document Name] Drawing

[Figure 1]

The nucleotide sequences of C2 region in  
subtypes of HIV-1

|           |  |
|-----------|--|
| SUBTYPE A | TGTAataccTCagccattTAcAcAgGctTgtCCaAaggTatCCTTTGAgCCaATTCCCATA      |
| SUBTYPE B | -----c-----t-----c-----  |
| SUBTYPE C | -----A-----a-----C-----c-----t-----c-----T-----                    |
| SUBTYPE D | -----g-----a-----  |
| SUBTYPE E | -----T-----T-----aG-----a-----T-----t-----                         |
| SUBTYPE F | -----A-----T-----GG-----T-----                                     |
| SUBTYPE G | -----gt-----A-----A-----ga?T-----c-----                            |
| SUBTYPE H | -----GT-----A-----GAGT-----T-----                                  |
| SUBTYPE A | caTTATTGtgCccCaGCTGGtTttGCgATtCTAAagTGtAa?gataaggagTTcaatGGA       |
| SUBTYPE B | -----g-----t-----a-----  |
| SUBTYPE C | -----t-----a-----ta-----aca-----g                                  |
| SUBTYPE D | -----a-----a-----a-----A-----g                                     |
| SUBTYPE E | -----a-----t-----a-----t-----T-----a-----t-----g                   |
| SUBTYPE F | -----T-----A-----T-----aA-----G                                    |
| SUBTYPE G | -----T-----t-----gg-----a-----?                                    |
| SUBTYPE H | -----T-----G-----A-----GG-----A-----G                              |
| SUBTYPE A | acAGGgccatGcaagAATGTcAGcaCaGTaCAATGcACacATGGaATcAagCCAGtagTa       |
| SUBTYPE B | -----a-----t-----ca-----t-----t-----g-----                         |
| SUBTYPE C | -----a-----c-----t-----t-----t-----g-----                          |
| SUBTYPE D | -----?-----a-----t-----g-----t-----g-----g                         |
| SUBTYPE E | -----t-----A-----T-----T-----G-----                                |
| SUBTYPE F | -----g-----T-----T-----A-----g-----                                |
| SUBTYPE G | -----A-----T-----a-----T-----T-----T-----g-----                    |
| SUBTYPE H | -----G-----AA-----T-----A-----T-----T-----G-----                   |
| SUBTYPE A | tCAACTCAaCTgcTGtTaAATGGcAGtctTAGCagaAgaa???gaggTAatgaTtagaTCT      |
| SUBTYPE B | -----a-----t-----c-----...-----g-----a-----                        |
| SUBTYPE C | -----?-----g-----...-----a-----A-----                              |
| SUBTYPE D | -----t-----T-----C-----...-----A-----A-----c-----                  |
| SUBTYPE E | -----t-----a-----c-----g-----t-----...-----ta-----A-----c-----     |
| SUBTYPE F | -----T-----A-----C-----GTCAAATG-CAGTTT-C-----?a-----a-----         |
| SUBTYPE G | -----gAaaataTcacAaAcAATgccaaaaAccaTAaTaGTacAgcTtg??aagcctGTAA?aATt |
| SUBTYPE B | -----t-----gg-----t-----gaa-----g-----at-----ga-----               |
| SUBTYPE C | -----c-----g-----t-----a-----t-----aAtg-----at-----ga-----         |
| SUBTYPE D | -----c-----t-----t-----?-----AATG-----t-----?c-----                |
| SUBTYPE E | -----C-----G-----C-----AAT-----At-----Ga-----C-----                |
| SUBTYPE F | -----c-----t-----g-----t-----A-----?-----AATg-----At-----ca-----   |
| SUBTYPE G | -----c?-----g-----gt-----g-----AAt-----a?-----a-----ga-----        |
| SUBTYPE H | -----c-----g-----a-----gt-----AAt-----a?-----g-----                |
| SUBTYPE A | aatTGT   |
| SUBTYPE B | -----  |
| SUBTYPE C | gtg---   |
| SUBTYPE D | -----  |
| SUBTYPE E | -----  |
| SUBTYPE F | -----  |
| SUBTYPE G | ??---  |
| SUBTYPE H | --?---   |




[Figure 2]


The nucleotide sequences of C3 region in  
subtypes of HIV-1

|           |   |
|-----------|---|
| SUBTYPE A | TGTAatgTcAgtaga?cagaaTGGaAtaaaactttTacaa?aggtagcta?acAatTAaga |
| SUBTYPE B | ----ca-t-----g-a-----c-----a-c-a--t---A-----                  |
| SUBTYPE C | ----cA-T---a-ga?a-----?-----a-----ag-a-a-----gc-              |
| SUBTYPE D | ----a-T---a-ga?a-----c-----a-a-----g-                         |
| SUBTYPE E | ---G-gA-T-A-g-A-a-----g-g-----a-c-----a-ga-a-----a-           |
| SUBTYPE F | ----c-t---g-a--C-----?---g-a-----a?ggc-a-g---ag               |
| SUBTYPE G | -----t-----a?-a-t-----?g-g-tG-----ga-t--?a??gc?--C--a-        |
| SUBTYPE H | -----T-----g?-a-?-----g-g-tg--?-?a-----?-?c-----?a-           |
|           |   |
| SUBTYPE A | aaa.....tacTtt?????????aaacaaaaca...??????ataatcTTtgctaac...  |
| SUBTYPE B | g--??????c-a---...g-g-----t-----g-----aa-c-a???               |
| SUBTYPE C | g--.....c---ccct-----T-----aa-----acca...                     |
| SUBTYPE D | g-c?.....cTtc-----aca-----t---aaacCa...                       |
| SUBTYPE E | g-g.....C-----a-t--T-G-----caaCCA???                          |
| SUBTYPE F | tct.....c-t--c-----tgC-----aa--aactcA...                      |
| SUBTYPE G | g--.....at-----?c???-----c---aaCtCA...                        |
| SUBTYPE H | ---.....?-----a---t---??.....c---aaacca???                    |
|           |   |
| SUBTYPE A | ?cctcaGGaGGGGAT?TaGAAaTtacAACacAtAgttTTAaTTGTggAgGagaattttTtc |
| SUBTYPE B | t-----cCc-----gt--tg--c-----g-----                            |
| SUBTYPE C | t-----cc-----c-----a-----                                     |
| SUBTYPE D | t-----ccc-----c-----g-----                                    |
| SUBTYPE E | c-----a--C-----tg--ca-----A--g-----                           |
| SUBTYPE F | t-----CC-----tg-----a-----                                    |
| SUBTYPE G | t-tg-----cC-----a-----  |
| SUBTYPE H | t-----Cc-----?-----a-----                                     |
|           |   |
| SUBTYPE A | TAtTGc  |
| SUBTYPE B | --c--t  |
| SUBTYPE C | -----   |
| SUBTYPE D | --C---  |
| SUBTYPE E | -----   |
| SUBTYPE F | --C---  |
| SUBTYPE G | -----t  |
| SUBTYPE H | -----t  |

[Figure 3]



[Figure 4]



[Document Name] Abstract

[Abstract]

[Object] A method for determining HIV-1 subtypes, which comprises the steps of amplifying nucleic acid using as a target sequence a portion of a nucleotide sequence of the env gene of HIV-1 wherein at least one of the 5' terminal and 3' terminal nucleotide sequences is different depending on the HIV-1 subtype, and detecting the subtype depending on whether or not the nucleic acid has been amplified. A kit for determining HIV-1 subtypes, comprising a primer pair in which a target sequence is a portion of a nucleotide sequence of the env gene of HIV-1 wherein at least one of the 5' terminal and 3' terminal nucleotide sequences is different depending on the subtype.

[Method for Achieving the Object] The present invention provides a simple method for determining HIV-1 subtypes. The present invention also provides an effective method for determining HIV-1 subtypes.

[Selected Figure] None